

DPNU-94-42
SNUTP-94-98
hep-th@xxx/9410119

Comment on “Obstructions to the Existence of Static Cosmic Strings in an Abelian Higgs Model”

Chanju Kim*

*Center for Theoretical Physics and Department of Physics
Seoul National University, Seoul 151-742, Korea*

Yoonbai Kim†

*Department of Physics, Nagoya University
Nagoya 464-01, Japan*

There has been considerable interest in the possible cosmic string configurations and Y. Yang has recently derived intriguing obstructions to the existence of cylindrically symmetric solutions in the Bogomol'nyi critical phase of the Einstein-Maxwell-Higgs system [1]. In this comment, we would like to mention some earlier results and recent progress [2, 3] which were not reflected in Ref. [1] but are important in connection with the obstructions he claimed. To our opinion, some of his obstructions are not consistent with the results of Refs. [2, 3] which we briefly summarize below. Also, we give a new condition stronger than those found before.

In the self-dual Abelian Higgs model coupled with gravity, once we restrict our interest to the string configurations parallel to the z axis and satisfying Bogomol'nyi equations, Einstein equations turn out to be solved under the general stationary metric and the whole solutions can be described by a single second-order equation for the amplitude of the Higgs field [4]

$$\begin{aligned} \partial^2 \ln \frac{f^2}{\prod_{p=1}^n |z - z_p|^2} \\ = e^{h+\bar{h}} \left(\frac{f^2 e^{-(f^2-1)}}{\prod_{p=1}^n |z - z_p|^2} \right)^{4\pi G v^2} (f^2 - 1), \end{aligned} \quad (1)$$

where f is the dimensionless scalar amplitude rescaled by the vacuum expectation value v , $h(z)$ ($\bar{h}(\bar{z})$) is a holomorphic (antiholomorphic) function which reduces to a constant, say F , in the cylindrically symmetric case, and z_p represents the center of p -th vortex. [The gravitational constant G in our notation corresponds to $G/2$ of Ref. [1].]

The finite energy condition for f is not simply that of flat case, $f(\infty) = 1$, but more complicated. For cylindrically symmetric case, it is given by

$$f(\infty) = \begin{cases} 1 \text{ or } 0 & \text{if } 0 < 4\pi G v^2 n \leq 1 \\ f_\infty, & 0 \leq f_\infty \leq 1 \text{ if } 4\pi G v^2 n > 1, \end{cases} \quad (2)$$

where n is the vorticity of the vortices superimposed at the origin (i.e., $f \sim f_0 r^n$ near the origin). Other than the well-known conic solutions which corresponds to $f(\infty) = 1$ with

$0 < 4\pi G v^2 n < 1$, non-conic solutions were first found by Linet [2] for $4\pi G v^2 n = 1$. In this case Eq. (1) (for the cylindrically symmetric case) is integrated to the first-order equation

$$\frac{df}{dr} = \frac{f}{r} \sqrt{n^2 - \frac{Fn}{2} f^{\frac{2}{n}} e^{-(f^2-1)}}. \quad (3)$$

Based on this, one can obtain the solutions for which the underlying two-manifold Σ is an (noncompact) asymptotic cylinder if $f(\infty) = 1$ with $F = 2n$ and $\Sigma \approx S^2$ (compact) if $f(\infty) = 0$ with $n = 1$ and $F > 2$. Here, the condition $n = 1$ which was overlooked in Refs. [2, 3] is necessary for the regularity of the solutions on the whole manifold $\Sigma \approx S^2$ and hence there exist only $N = 2$ solutions on S^2 . [One may easily convince himself that, otherwise, the phase of the Higgs field would not be regular everywhere outside the vortex points.] The maximum circumference of the cylinder and that of two-sphere along the tropical line has a fixed value $\sqrt{32\pi^3 G} n/e$ independently of F [3]. There is no other type of regular solutions to Eq. (1) in a geodesically complete manifold [3]. [In fact, in Ref. [3], we considered more general system which may have external point particles.] That is, among the many possibilities in Eq. (2), regular finite-energy solutions exist only for the aforementioned three cases.

From these, we see that (i) in Ref. [1], the second condition of Eq. (7) in Theorem 2 and Eq. (14) in Theorem 5 (and subsequent discussions) can not be true since there exist asymptotic-cylinder solutions; (ii) contrary to Yang's claim of the existence of $|N| \geq 3$ solutions for $\Sigma \approx S^2$, regular solutions exist only for $N = 2$. Correspondingly, in this case, vacuum expectation value v is quantized uniquely as $v = 1/\sqrt{4\pi G}$ rather than v can take arbitrary value of Eq. (4) of Ref. [1].

We thank Choonkyu Lee for reading the manuscript. This work was supported by the Korea Science and Engineering Foundation through Center for Theoretical Physics, Seoul National University and JSPS under #93033 (Y.K.).

References

* E-mail address: cjkim@phyb.snu.ac.kr

† E-mail address: yoonbai@eken.phys.nagoya-u.ac.jp

- [1] Y. Yang, Phys. Rev. Lett. **73**, 10 (1994).
- [2] B. Linet, Class. Quantum Grav. **7** (1990) L79.
- [3] C. Kim, and Y. Kim, Phys. Rev. **50**, 1040 (1994).
- [4] B. Linet, Gen. Rel. Grav. **20** (1988) 451; A. Comtet, and G. W. Gibbons, Nucl. Phys. B299 (1988) 719.